

1. A red die and a blue die are rolled. Let  $A$  be the event that the red die rolls 2, let  $B$  be the event that the sum of the rolls is 5, and let  $C$  be the event that the sum of the rolls is 7. Are  $A$  and  $B$  independent events? How about  $A$  and  $C$ ?

$$\text{First, } P(A) = \frac{1}{6} \quad \text{and} \quad P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{36}}{\frac{4}{36}} = \frac{1}{4}.$$

Since  $P(A) \neq P(A|B)$ , events  $A$  and  $B$  are dependent.

$$\text{Second, } P(A|C) = \frac{P(A \cap C)}{P(C)} = \frac{\frac{1}{36}}{\frac{6}{36}} = \frac{1}{6}.$$

Since  $P(A) = P(A|C)$ , events  $A$  and  $C$  are independent.

2. A sequence of  $n$  independent trials are to be performed. Each trial results in a success with probability  $p$  and a failure with probability  $1 - p$ . What is the probability that...

- (a) ...all trials result in successes?

Let  $A_i$  be the event that trial  $i$  results in success.

$$\text{By independence, } P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1)P(A_2)\dots P(A_n) = \underbrace{p \cdot p \cdot \dots \cdot p}_{n \text{ factors}} = p^n$$

- (b) ...at least one trials results in a success?

$$\text{Probability of no successes: } P(A'_1 \cap A'_2 \cap \dots \cap A'_n) = (1-p)^n$$

$$\text{Probability of at least one success: } 1 - P(A'_1 \cap \dots \cap A'_n) = 1 - (1-p)^n$$

- (c) ...exactly  $k$  trials result in successes?

We will see this again in Chapter 2.

Any particular sequence of  $k$  successes and  $n-k$  failures occurs with probability  $p^k (1-p)^{n-k}$ . There are  $\binom{n}{k}$  such sequences.

$$\text{Thus, } P(\text{exactly } k \text{ successes}) = \binom{n}{k} p^k (1-p)^{n-k}.$$

↑  
This factor is important!

3. Consider an urn containing four balls, numbered 110, 101, 011, and 000. One ball is drawn at random. For  $k = 1, 2, 3$ , let  $A_k$  be the event that the  $k^{\text{th}}$  digit is a 1 on the ball that is drawn.

- (a) Are the events  $A_1$ ,  $A_2$ , and  $A_3$  pairwise independent? Why or why not?

$$\text{Yes: } P(A_i) = \frac{1}{2} \quad \text{for any } i \in \{1, 2, 3\}$$

$$P(A_i \cap A_j) = \frac{1}{4} \quad \text{for any } i, j \in \{1, 2, 3\}$$

(b) Are the events  $A_1, A_2,$  and  $A_3$  mutually independent? Why or why not?

No:  $P(A_1 \cap A_2 \cap A_3) = 0 \neq P(A_1)P(A_2)P(A_3)$

4. If  $A$  and  $B$  are independent events with positive probability, show that they cannot be mutually exclusive.


Given:  $P(A) > 0, P(B) > 0,$  and  $P(A \cap B) = P(A)P(B).$

Thus  $P(A \cap B) > 0,$  so the events can occur simultaneously, meaning they are not mutually exclusive.

5. Create an example of three events  $A, B, C$  such that  $P(A \cap B \cap C) = P(A)P(B)P(C)$  but the events are not mutually independent.

This is tricky! Trial-and-error is a fine approach.

Here is one example:



$$P(A) = P(B) = P(C) = \frac{1}{2}$$

$$P(A \cap B \cap C) = \frac{1}{8} = P(A)P(B)P(C)$$

However:

$$P(A \cap B) = \frac{9}{40} \neq P(A)P(B)$$

(and similarly for  $A \cap C$  and  $B \cap C$ )

$A, B,$  and  $C$  are not pairwise independent, and thus not mutually independent.